

HUREWICZ SETS OF REALS WITHOUT PERFECT SUBSETS

DUŠAN REPOVŠ, BOAZ TSABAN, AND LYUBOMYR ZDOMSKYY

ABSTRACT. We show that even for subsets X of the real line which do not contain perfect sets, the Hurewicz property does not imply the property $S_1(\Gamma, \Gamma)$, asserting that for each countable family of open γ -covers of X , there is a choice function whose image is a γ -cover of X . This settles a problem of Just, Miller, Scheepers, and Szeptycki. Our main result also answers a question of Bartoszyński and the second author, and implies that for $C_p(X)$, the conjunction of Sakai's strong countable fan tightness and the Reznichenko property does not imply Arhangel'skii's property α_2 .

1. INTRODUCTION

By a *set of reals* we mean a separable, zero-dimensional, and metrizable space (such spaces are homeomorphic to subsets of the real line $\mathbb{R}$). Fix a set of reals X . Let $\mathcal{O}$ denote the collection of all open covers of X . An open cover $\mathcal{U}$ of X is a γ -cover of X if it is infinite and for each $x \in X$, x is a member of all but finitely many members of $\mathcal{U}$. Let Γ denote the collection of all open γ -covers of X . Motivated by Menger's work, Hurewicz [6] introduced the *Hurewicz property* $U_{\text{fin}}(\mathcal{O}, \Gamma)$:

For each sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of members of $\mathcal{O}$ which do not contain a finite subcover, there exist finite sets $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \mathbb{N}$, such that $\{\cup \mathcal{F}_n : n \in \mathbb{N}\} \in \Gamma$.

Every σ -compact space satisfies $U_{\text{fin}}(\mathcal{O}, \Gamma)$, but the converse fails [7, 2].

Let $\mathcal{A}$ and $\mathcal{B}$ be any two families. Motivated by works of Rothberger, Scheepers introduced the following prototype of properties [12].

$S_1(\mathcal{A}, \mathcal{B})$: For each sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of members of $\mathcal{A}$, there exist members $U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$, such that $\{U_n : n \in \mathbb{N}\} \in \mathcal{B}$.

It is easy to see that $U_{\text{fin}}(\mathcal{O}, \Gamma) = U_{\text{fin}}(\Gamma, \Gamma)$, and therefore $S_1(\Gamma, \Gamma)$ implies $U_{\text{fin}}(\mathcal{O}, \Gamma)$ [12]. However, a set of reals satisfying $S_1(\Gamma, \Gamma)$ cannot contain perfect subsets [7]. It follows that, for example, $\mathbb{R}$ satisfies $U_{\text{fin}}(\mathcal{O}, \Gamma)$ but not $S_1(\Gamma, \Gamma)$. In the fundamental paper [7], we are asked whether there are *nontrivial* examples showing that $U_{\text{fin}}(\mathcal{O}, \Gamma)$ does not imply $S_1(\Gamma, \Gamma)$.

Problem 1.1 (Just, Miller, Scheepers, Szeptycki [7]). *Let X be a set of reals which does not contain a perfect set, but which does have the Hurewicz property. Does X then satisfy $S_1(\Gamma, \Gamma)$?*

We give a negative answer that also yields a new result concerning function spaces.

1991 *Mathematics Subject Classification*. Primary: 37F20; Secondary 26A03, 03E75.

The first and the third authors were supported by the Slovenian Research Agency grants P1-0292-0101-04 and BI-UA/04-06-007. The second author was partially supported by the Koshland Center for Basic Research.

2. THE MAIN THEOREM

We prove a stronger assertion than what is needed to settle Problem 1.1; this will be useful for the next section. Let C_Γ denote the collection of all *clopen* γ -covers of X . Clearly, $S_1(\Gamma, \Gamma)$ implies $S_1(C_\Gamma, C_\Gamma)$.¹ The hypothesis in the following theorem is a consequence of the Continuum Hypothesis. See [3] for a survey of the involved cardinals.

Theorem 2.1. *Assume that $\mathfrak{b} = \mathfrak{c}$. There exists a set of reals X such that:*

- (1) *X does not contain a perfect set;*
- (2) *All finite powers of X have the Hurewicz property $U_{\text{fin}}(\mathcal{O}, \Gamma)$; and*
- (3) *No set of reals containing X satisfies $S_1(C_\Gamma, C_\Gamma)$.*

Theorem 2.1 is proved in three steps. The first step is analogous to Theorem 4.2 of [5], and will be used to show that the constructed set is not contained in a set of reals satisfying $S_1(C_\Gamma, C_\Gamma)$. Say that a convergent sequence $\{x_n\}_{n \in \mathbb{N}}$ is *nontrivial* if $\lim_n x_n \notin \{x_n : n \in \mathbb{N}\}$.

Lemma 2.2. *Let X be a subspace of a zero-dimensional metrizable space Y satisfying $S_1(C_\Gamma, C_\Gamma)$, and $\{x_n^m\}_{n \in \mathbb{N}}$, $m \in \mathbb{N}$, be nontrivial convergent sequences in X . Then: There are a countable closed cover $\{F_k : k \in \mathbb{N}\}$ of X and an infinite $A \subseteq \mathbb{N}$, such that $F_k \cap \{x_n^m : n \in A\}$ is finite for all k, m .*

Proof. Let d be a metric on Y which generates its topology. For each m , do the following. Let $x_m = \lim_n x_n^m$, and for each n take a clopen neighborhood C_n^m of x_n^m in Y , whose diameter is smaller than $d(x_n^m, x_m)/2$. For each m, n , set

$$U_n^m = Y \setminus (C_n^0 \cup C_n^1 \cup \dots \cup C_n^m).$$

For each m , $\{U_n^m : n \in \mathbb{N}\}$ is a clopen γ -cover of Y . Apply $S_1(C_\Gamma, C_\Gamma)$ to get $f \in \mathbb{N}^\mathbb{N}$ such that $\{U_{f(m)}^m : m \in \mathbb{N}\}$ is a (clopen) γ -cover of Y . As $U_{f(m)}^m \subseteq Y \setminus C_{f(m)}^0$ for each m , we have that the image A of f is infinite.

For each k , let $F_k = \bigcap_{i \geq k} U_{f(i)}^i$. $\{F_k : k \in \mathbb{N}\}$ is a closed (γ -)cover of Y . Fix k and m . If n is large enough and $n \in A$, then $n = f(i)$ with $i \geq m, k$. As $x_n^m = x_{f(i)}^m \in C_{f(i)}^m$ and $i \geq m$, $x_n^m \notin U_{f(i)}^i$. As $i \geq k$, $U_{f(i)}^i \supseteq F_k$, and therefore $x_n^m \notin F_k$. $\square$

To make sure that our constructed set does not contain a perfect set and that it satisfies the Hurewicz property in all finite powers, we will use the following. Let $\overline{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$ be the one point compactification of $\mathbb{N}$, and $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$ be the collection of all nondecreasing elements f of $\overline{\mathbb{N}}^\mathbb{N}$ (endowed with the Tychonoff product topology) such that $f(n) < f(n+1)$ whenever $f(n) < \infty$. $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$ is homeomorphic to the Cantor space (see [13] for an explicit homeomorphism), and can therefore be viewed as a set of reals.

Let S be the family of all nondecreasing finite sequences in $\mathbb{N}$. For $s \in S$, $|s|$ denotes its length. For each $s \in S$, define $q_s \in \overline{\mathbb{N}}^{\uparrow \mathbb{N}}$ by $q_s(n) = s(n)$ if $n < |s|$, and $q_s(n) = \infty$ otherwise. Let Q be the collection of all these elements q_s . Q is dense in $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$.

¹It is an open problem whether the converse implication holds [4, 11].

For a set D and $f, g \in \mathbb{N}^D$, $f \leq^* g$ means: $f(d) \leq g(d)$ for all but finitely many $d \in D$. A $\mathfrak{b}$ -scale is an unbounded (with respect to $\leq^*$) set $\{f_\alpha : \alpha < \mathfrak{b}\} \subseteq \mathbb{N}^{\mathbb{N}}$ of increasing functions, such that $f_\alpha \leq^* f_\beta$ whenever $\alpha < \beta$.

Theorem 2.3 (Bartoszyński-Tsaban [2]). *Let $X \subseteq \overline{\mathbb{N}}^{\uparrow \mathbb{N}}$ be a union of a $\mathfrak{b}$ -scale and Q . Then X contains no perfect subset, and all finite powers of H satisfy the Hurewicz property $\mathcal{U}_{\text{fin}}(\mathcal{O}, \Gamma)$.*

For each $s \in S$, $\{q_{s^\frown n}\}_{n \in \mathbb{N}}$ (where $\frown$ denotes a concatenation of sequences) is a nontrivial convergent sequence in $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$,² and

$$\lim_{n \rightarrow \infty} q_{s^\frown n} = q_s.$$

The following will be used in our construction.

Lemma 2.4. *Let X be a closed subspace of $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$. If $X \cap \{q_{s^\frown n} : n \in \mathbb{N}\}$ is finite for each $s \in S$, then there exists $\phi : S \rightarrow \mathbb{N}$ such that for all $x \in X$ and all $n \geq 2$, $x(n) \geq \phi(x \upharpoonright n)$ implies $x(n+1) \leq \phi(x \upharpoonright (n+1))$.*

Proof. For each $s \in S$, let $k(s)$ be such that $q_{s^\frown k} \in \overline{\mathbb{N}}^{\uparrow \mathbb{N}} \setminus X$ for all $k \geq k(s)$. As X is closed in $\overline{\mathbb{N}}^{\uparrow \mathbb{N}}$, for each $k \geq k(s)$ there is $m(s, k)$ such that

$$\{z \in \overline{\mathbb{N}}^{\uparrow \mathbb{N}} : z \upharpoonright (|s| + 1) = s^\frown k, z(|s| + 1) > m(s, k)\} \cap X = \emptyset.$$

(Note that $\{z \in \overline{\mathbb{N}}^{\uparrow \mathbb{N}} : z \upharpoonright (|s| + 1) = s^\frown k, z(|s| + 1) > m\}$, $m \in \mathbb{N}$, is a neighborhood base at $q_{s^\frown k}$.) Define $\phi : S \rightarrow \mathbb{N}$ by

$$\phi(s) = \max\{k(s), m(s \upharpoonright (|s| - 1)), s(|s| - 1)\}$$

when $|s| \geq 2$, and by $\phi(s) = 0$ when $|s| < 2$. Let $x \in X$ and $n \geq 2$. If $x(n) \geq \phi(x \upharpoonright n)$, then $x(n) \geq k(x \upharpoonright n)$, and hence $x(n+1) \leq m(x \upharpoonright n, x(n)) \leq \phi(x \upharpoonright (n+1))$. $\square$

It remains to prove the following.

Proposition 2.5. *Assume that $\mathfrak{b} = \mathfrak{c}$. There exists a $\mathfrak{b}$ -scale $B = \{b_\alpha : \alpha < \mathfrak{b}\}$ such that for each closed cover $\{F_n : n \in \mathbb{N}\}$ of $B \cup Q$ and each infinite set $A \subseteq \mathbb{N}$, there are n and $s \in S$ such that $F_n \cap \{q_{s^\frown k} : k \in A\}$ is infinite.*

Proof. Let $\{A_\alpha : \alpha < \mathfrak{c}\}$ be an enumeration of all infinite subsets of $\mathbb{N}$, such that for each infinite $A \subseteq \mathbb{N}$, there are $\mathfrak{c}$ many $\alpha < \mathfrak{c}$ with $A_\alpha = A$.

As $\mathfrak{b} = \mathfrak{d} = \mathfrak{c}$, there is a (standard) scale in $\mathbb{N}^S$, that is, a family $\{\phi_\alpha : \alpha < \mathfrak{c}\} \subseteq \mathbb{N}^S$ such that:

- (1) For each $\phi \in \mathbb{N}^S$, there is $\beta < \mathfrak{c}$ such that $\phi \leq^* \phi_\beta$; and
- (2) For all $\alpha < \beta < \mathfrak{c}$, $\phi_\alpha \leq^* \phi_\beta$.

For an infinite $A \subseteq \mathbb{N}$, let $\overline{A} = A \cup \{\infty\}$, and

$$\overline{A}^{\uparrow \mathbb{N}} = \{x \in \overline{\mathbb{N}}^{\uparrow \mathbb{N}} : x(n) \in \overline{A} \text{ for all } n\}$$

The order isomorphism between $A \cup \{\infty\}$ and $\mathbb{N} \cup \{\infty\}$ induces an order isomorphism $\Psi_A : \overline{A}^{\uparrow \mathbb{N}} \rightarrow \overline{\mathbb{N}}^{\uparrow \mathbb{N}}$.

²Strictly speaking, $q_{s^\frown n} \notin \overline{\mathbb{N}}^{\uparrow \mathbb{N}}$ when $n < s(|s| - 1)$, but since we are dealing with convergent sequences, we can ignore the first few elements.

By induction on $\alpha < \mathfrak{b} = \mathfrak{c}$, construct a $\mathfrak{b}$ -scale $B = \{b_\alpha : \alpha < \mathfrak{c}\}$ such that for each $\alpha < \mathfrak{c}$, $b_\alpha \in (A_\alpha)^{\uparrow \mathbb{N}}$, and

$$\Psi_{A_\alpha}(b_\alpha)(n) > \phi_\alpha(\Psi_{A_\alpha}(b_\alpha) \upharpoonright n)$$

for all $n \geq 2$.

We claim that $X = B \cup Q$ is as required. Indeed, let A be an infinite subset of $\mathbb{N}$. Take an increasing enumeration $\{\beta_\alpha : \alpha < \mathfrak{c}\}$ of $\{\alpha < \mathfrak{c} : A_\alpha = A\}$. For each $\alpha < \mathfrak{c}$, $b_{\beta_\alpha} \in \overline{A}^{\uparrow \mathbb{N}}$. Set $c_\alpha = \Psi_A(b_{\beta_\alpha})$, and $C = \{c_\alpha : \alpha < \mathfrak{c}\}$. By the construction of the functions b_α ,

$$c_\alpha(n) > \phi_{\beta_\alpha}(c_\alpha \upharpoonright n) \geq \phi_\alpha(c_\alpha \upharpoonright n)$$

for all but finitely many n .

Let $\{K_m : m \in \mathbb{N}\}$ be a closed cover of $C \cup Q$. Then there are m and $s \in S$ such that $K_m \cap \{q_{s \cdot k} : k \in \mathbb{N}\}$ is infinite. Otherwise, by Lemma 2.4, for each m there is $\psi_m \in \mathbb{N}^S$ such that for all $x \in K_m$ and $n \geq 2$, $x(n) \geq \psi_m(x \upharpoonright n)$ implies $x(n+1) \leq \psi_m(x \upharpoonright (n+1))$. Let $\alpha < \mathfrak{c}$ be such that for each m , $\phi_\alpha(s) \geq \psi_m(s)$ for all but finitely many $s \in S$. It is easy to verify that $c_\alpha \notin K_m$ for all m ; a contradiction.

Now consider any closed cover $\{F_m : m \in \mathbb{N}\}$ of $B \cup Q$ and set $K_m = \Psi_A(F_m \cap \overline{A}^{\uparrow \mathbb{N}})$. Let $s \in S$ and m be such that $K_m \cap \{q_{s \cdot k} : k \in \mathbb{N}\}$ is infinite. Then for $\tilde{s} \in S$ such that $\tilde{s}(i)$ is the $s(i)$ 'th element of A for each $i < |s|$, we have that $F_m \cap \{q_{\tilde{s} \cdot k} : k \in A\}$ is infinite. $\square$

This completes the proof of Theorem 2.1. The following corollary of Theorem 2.1 answers in the negative Problem 15(1) of Bartoszyński and the second author [2].

Corollary 2.6. *The union of a $\mathfrak{b}$ -scale and Q need not satisfy $S_1(\Gamma, \Gamma)$.* $\square$

3. REFORMULATION FOR SPACES OF CONTINUOUS FUNCTIONS

Let Y be a (not necessarily metrizable) topological space. For $y \in Y$ and $A \subseteq Y$, write $\lim A = y$ if A is countable, and an (any) enumeration of A converges nontrivially to y . Let $\Gamma_y = \{A \subseteq Y : \lim A = y\}$. Y has the Arhangel'skiĭ *property* α_2 [1] if $S_1(\Gamma_y, \Gamma_y)$ holds for all $y \in Y$.

Fix a set of reals X . $C_p(X)$ is the subspace of the Tychonoff product $\mathbb{R}^X$ consisting of the continuous functions. It was recently discovered, independently by Bukovský-Haleš [4] and by Sakai [11], that $C_p(X)$ has the property α_2 if, and only if, X satisfies $S_1(C_\Gamma, C_\Gamma)$.

Many additional connections of this type are studied in the literature. For families $\mathcal{A}$ and $\mathcal{B}$, consider the following prototype [12].

$S_{\text{fin}}(\mathcal{A}, \mathcal{B})$: For each sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of members of $\mathcal{A}$, there exist finite subsets $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \mathbb{N}$, such that $\bigcup_n \mathcal{F}_n \in \mathcal{B}$.

For a topological space Y and $y \in Y$, let $\Omega_y = \{A \subseteq Y : y \in \overline{A} \setminus A\}$. Y has the Arhangel'skiĭ *countable fan tightness* [1] if $S_{\text{fin}}(\Omega_y, \Omega_y)$ holds for each $y \in Y$. Y has the *Reznichenko property* if for each $y \in Y$ and each $A \in \Omega_y$, there are pairwise disjoint finite sets $F_n \subseteq A$, $n \in \mathbb{N}$, such that each neighborhood U of y intersects F_n for all but finitely many n .

For sets of reals X , $C_p(X)$ has countable fan tightness and the Reznichenko property if, and only if, all finite powers of X have the Hurewicz property $U_{\text{fin}}(\mathcal{O}, \Gamma)$ [9]. Thus, Theorem 2.1 can be reformulated as follows.

Theorem 3.1. *Assume that $\mathfrak{b} = \mathfrak{c}$. There exists a set of reals X without perfect subsets, such that $C_p(X)$ has countable fan tightness and the Reznichenko property, but does not have the Arhangel'skiĭ property α_2 .* $\square$

A topological space Y has the *Sakai strong countable fan tightness* if $S_1(\Omega_y, \Omega_y)$ holds for each $y \in Y$. Sakai proved that for sets of reals, $C_p(X)$ has strong countable fan tightness if, and only if, all finite powers of X satisfy $S_1(\mathcal{O}, \mathcal{O})$ [10]. For sets of reals X , $C_p(X)$ has strong countable fan tightness and the Reznichenko property if, and only if, all finite powers of X satisfy $U_{\text{fin}}(\mathcal{O}, \Gamma)$ as well as $S_1(\mathcal{O}, \mathcal{O})$ [8].

If $\mathfrak{b} \leq \text{cov}(\mathcal{M})$ and X is a union of a $\mathfrak{b}$ -scale and Q , then all finite powers of X satisfy $U_{\text{fin}}(\mathcal{O}, \Gamma)$ as well as $S_1(\mathcal{O}, \mathcal{O})$ [2]. As the Continuum Hypothesis (or just Martin's Axiom) implies that $\mathfrak{b} = \text{cov}(\mathcal{M}) = \mathfrak{c}$, we have the following.

Corollary 3.2. *Even for $C_p(X)$ where X is a set of reals, the conjunction of strong countable fan tightness and the Reznichenko property does not imply the Arhangel'skiĭ property α_2 .* $\square$

4. CONCLUDING REMARKS AND OPEN PROBLEMS

Our results are consistency results. What is not settled is whether the answers to the problems addressed in this paper are undecidable.

Problem 4.1. *Is it consistent that all sets of reals which have the Hurewicz property $U_{\text{fin}}(\mathcal{O}, \Gamma)$ but have no perfect subsets satisfy $S_1(\Gamma, \Gamma)$?*

Problem 4.2. *Is it consistent that each union of a $\mathfrak{b}$ -scale and Q satisfies:*

- (1) $S_1(\Gamma, \Gamma)$?
- (2) $S_1(\Gamma, \Gamma)$ in all finite powers?

Problem 4.3. *Is it consistent that for each set of reals X , if $C_p(X)$ has both strong countable fan tightness and the Reznichenko property, then $C_p(X)$ has the Arhangel'skiĭ property α_2 ?*

REFERENCES

- [1] A. V. Arhangel'skiĭ, *The frequency spectrum of a topological space and the classification of spaces*, Soviet Mathematical Doklady **13** (1972), 1186–1189.
- [2] T. Bartoszyński and B. Tsaban, *Hereditary topological diagonalizations and the Menger-Hurewicz Conjectures*, Proceedings of the American Mathematical Society **134** (2006), 605–615.
- [3] A. R. Blass, *Combinatorial cardinal characteristics of the continuum*, in: **Handbook of Set Theory** (M. Foreman, A. Kanamori, and M. Magidor, eds.), Kluwer Academic Publishers, Dordrecht, to appear. <http://www.math.lsa.umich.edu/~ablass/hbk.pdf>
- [4] L. Bukovský and J. Haleš, *QN-space, wQN-space and covering properties*, Topology and its Applications **154** (2007), 848–858.
- [5] L. Bukovský, I. Reclaw, and M. Repický, *Spaces not distinguishing pointwise and quasinormal convergence of real functions*, Topology and its Applications **41** (1991), 25–41.
- [6] W. Hurewicz, *Über Folgen stetiger Funktionen*, Fundamenta Mathematicae **9** (1927), 193–204.
- [7] W. Just, A. W. Miller, M. Scheepers, and P. J. Szeptycki, *The combinatorics of open covers II*, Topology and its Applications **73** (1996), 241–266.
- [8] Lj. D.R. Kočinac and M. Scheepers, *Function spaces and a property of Reznichenko*, Topology and its Applications **123** (2002), 135–143.
- [9] Lj. D.R. Kočinac and M. Scheepers, *Combinatorics of open covers (VII): Groupability*, Fundamenta Mathematicae **179** (2003), 131–155.

- [10] M. Sakai, *Property C'' and function spaces*, Proceedings of the American Mathematical Society **104** (1988), 917–919.
- [11] M. Sakai, *The sequence selection properties of $C_p(X)$* , Topology and its Applications **154** (2007), 552–560.
- [12] M. Scheepers, *Combinatorics of open covers I: Ramsey theory*, Topology and its Applications **69** (1996), 31–62.
- [13] B. Tsaban and L. Zdomskyy, *Scales, fields, and a problem of Hurewicz*, Journal of the European Mathematical Society, to appear.

DUŠAN REPOVŠ, INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA, P.O.B. 2964, LJUBLJANA, SLOVENIJA 1001.

E-mail address: `dusan.repovs@guest.arnes.si`

BOAZ TSABAN, DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN 52900, ISRAEL; AND DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL.

E-mail address: `tsaban@math.biu.ac.il`

LYUBOMYR ZDOMSKYY, KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, WÄHRINGER STR. 25, A-1090 VIENNA, AUSTRIA.

E-mail address: `lzdomsky@gmail.com`